

50X1-HUM

Declassified in Part - Sanitized Copy Approved for Release 2012/07/12 :
CIA-RDP80S01540R005500040014-8

Page Denied

Declassified in Part - Sanitized Copy Approved for Release 2012/07/12 :
CIA-RDP80S01540R005500040014-8

INFORMATION REPORT

50X1-HUM

REPORT

DATE DISTR. 26 May 1954

NO. OF PAGES 1

REQUIREMENT NO. RD

REFERENCES

50X1-HUM

50X1-HUM

SECRET CONTROL--U. S. OFFICIALS 'ONLY

AUG 18 2 50 PM '54
OSI/IC

STATE	ARMY	NAVY	AIR	FBI	AEC	OCD	X	
-------	------	------	-----	-----	-----	-----	---	--

50X1-HUM

(Note: Washington Distribution Indicated By "X"; Field Distribution By "#".) Form No. 51-61, January 1953

SECRET U. S. OFFICIALS ONLY

50X1-HUM

East Germany

Research on Rare Earths at the Institute for Inorganic Chemistry, Leipzig.

extinction with liquid ammonia of

The project undertaken by Mr. Zott called for the separation of mixtures of neutral and basic nitrates, obtained from ~~various rare earth nitrates~~ the partial thermal decomposition of mixtures of rare-earth-nitrates. It was expected that the neutral nitrates would dissolve much better in ammonia than the basic nitrates would. Results were good for the rare earths lying far apart on the atomic weight scale, but for rare earths in close proximity to each other the atomic weight differences were only 1-3 units. Other preliminary experiments Mr. Zott attempted to utilize the basically low differences in ionic migration velocities of the rare earths to separate them. The results are differences of atomic weight from 3.5 to 4.7 units. Comparisons are hardly possible with other methods because Mr. Zott worked with unknown mixtures.

^{with Mr. Zott}
The research group working in this consisted of Dr. Heinz Holzapfel, Dipl.Chem. Hans-Joachim Jahn, cand.chem. Joachim Nassonne, and cand.chem. Gerhard Zimmermann.

Since no up-to-date apparatus was available, such as a Roentgen-spectrograph, Qu 24, [quartz spectrograph] or mirror monochromator, the necessary apparatus had to be constructed by adapting what was on hand or by building it from parts of discarded instruments.

The impurity of chemicals was another handicap. It was possible, although very time-consuming, to purify what chemicals were needed.

As far as literature is concerned: old "classic" handbooks were available for preliminary work. Soviet literature has been received more regularly for some time now, but ~~much~~ as our study of the rare earths will progress, the need for foreign literature will increase.

The work plan is now divided into three parts: 1) as mentioned above, the building of the necessary apparatus. 2) Work to acquaint the group with the classic methods of rare-earth-chemistry. a) Determination of atomic weights. b) Development of a preparation "Crystals from liquor of Yttrium material". (The material used to start the experiments was found to be a potassium-rare-earth bisulfate.) 3) Work designed to increase the knowledge of the subject and to further develop methods in the field of rare earths. a) critical testing of the methods for the development of certain preparations of greatest purity. b) Testing of Neunhoeffer's work on the detection by fluorescence of rare earths.

Dipl. Chemist Kurt Bernhardt worked on preliminary experiments on the dependence of pH of the precipitation of Yttrium with ~~alizarin~~ the sodium salt of alizarin sulfonic acid.

Dipl. Chem Hermann Scholer experimented with separation experiments with Lanthanum-Neodymium and Lanthanum-samarium mixtures.

50X1-HUM

East Germany

Research on Rare Earths at the Institute for Inorganic Chemistry, Leipzig.

50X1-HUM
The project undertaken by Mr. Zott called for the ~~partial thermic decomposition of~~ extraction with liquid ammonia of mixtures of neutral and basic nitrates, obtained from ~~rare earths nitrates~~ the partial thermic decomposition of mixtures of rare-earth-nitrates. It was expected that the neutral nitrates would dissolve much better in ammonia than the basic nitrates would. Results were good for the rare earths lying far apart on the atomic weight scale, but for rare earths in close proximity to ~~the compounds were only 1-2~~ each other the atomic weight differences were only 1-3 units. Other preliminary experiments Mr. Zott attempted to utilize the basically low differences in ionic migration velocities of the rare earths to separate them. The results are differences of atomic weight from 3.5 to 4.7 units. Comparisons are hardly possible with other methods, because Mr. Zott worked with ~~unknown~~ mixtures of unknown composition.

The research group working ~~in this~~ ^{with Mr. Zott} consisted of Dr. Heinz Holzapfel, Dipl.Chem. Hans-Joachim Jahn, cand.chem. Joachim Massonne, and cand.chem. Gerhard Zimmermann.

Since no up-to-date apparatus was available, such as a Roentgeno-spectrograph, Qu 24, [quartz spectrograph] or mirror monochromator, the necessary apparatus had to be constructed by adapting what was on hand or by building it from parts of discarded instruments.

The impurity of chemicals was another handicap. It was possible, although very time-consuming, to purify what chemicals were needed.

As far as literature is concerned: old "classic" handbooks were available for preliminary work. Soviet literature has been received more regularly for some time now, but, ~~so far~~ as our study of the rare earths will progress, the need for foreign literature will increase.

The work plan is now divided into three parts: 1) as mentioned above, the building of the necessary apparatus. 2) Work to acquaint the group with the classic methods of rare-earth-chemistry. a) Determination of atomic weights. b) Development of a preparation "Crystals from liquor of Yttrium material". (The material used to start the experiments was found to be a potassium-rare-earth bisulfate.) 3) Work designed to increase the knowledge of the subject, and to further develop methods, in the field of rare earths. a) critical testing of the methods for the ~~pre~~ development of cerium preparations of greatest purity. b) Testing of Neunhoeffer's work on the detection by fluorescence of rare earths.

Dipl. Chemist Kurt Bernhardt worked on preliminary experiments on the dependence ~~on~~ pH of the precipitation of Yttrium with ~~alizerin~~ the sodium salt of alizarin sulfonic acid. Dipl. Chem. Hermann Scheler experimented with separation experiments with lanthanum-Neodymium and Lanthanum-samarium mixtures.

50X1-HUM

Declassified in Part - Sanitized Copy Approved for Release 2012/07/12 : CIA-RDP80S01540R005500040014-8
50X1-HUM

Page Denied

Next 7 Page(s) In Document Denied